

REFERENCES

- Anonymous, 2008. The Gazette of the Democratic Socialist Republic of Sri Lanka (Extraordinary) No 1534/18: National Environmental (Protection and Quality) Regulations, No. 1 of 2008.
- APHA, 2017. Standard Methods for the Examination of Water and Wastewater. 23rd edition. American Public Health Association, American Water Worker Association, Water Pollution Control Federation, New York, USA.
- Barberio, A., 2013. Bioassays with plants in the monitoring of water quality. INTECH Open Access Publisher.
- Basavaraju, H. C., & Chandru, S. (2008). Impact of distillery spentwash on the nutrients of leaves vegetables: An investigation. Asian Journal of Chemistry, 20(7), 5301.
- Baskar, M., Kayalvizhi, C., & Bose, M. (2003). Eco-friendly utilisation of distillery effluent in agriculture—A review. Agricultural Reviews, 24(1), 16-30.
- Baskar, M., Saravanan, A., Chitra, L., Dhevagi, P., Lenin, R. D., Pandiyarajan, P., & Ambast, S. K. (2013). Ecofriendly utilization of distillery waste water in agriculture. AICRP on Management of Salt Affected Soils and Use of Saline Water in Agriculture, 1-50.
- Basso, H. A. L., & Lopes, M. (2009). Sugar cane juice and molasses, beet molasses and sweet sorghum: Composition and usage.
- Bharagava, R. N., & Chandra, R. (2010). Effect of bacteria treated and untreated post-methanated distillery effluent (PMDE) on seed germination, seedling growth and amylase activity in *Phaseolus mungo* L. Journal of Hazardous Materials, 180(1-3), 730-734.
- Carita, R. and Marin-Morales, M.A., 2008. Induction of chromosome aberrations in the *Allium cepa* test system caused by the exposure of seeds to industrial effluents contaminated with azo dyes. Chemosphere, 72(5), pp.722-725.

Chandra, R., Bharagava, R. N., & Rai, V. (2008). Melanoidins as major colourant in sugarcane molasses based distillery effluent and its degradation. *Bioresource technology*, 99(11), 4648-4660.

Chidankumar, C. S., & Chandraju, S. (2009). Impact of Distillery Spentwash Irrigation on Yield of Some Root Vegetables. *Nature Environment and Pollution Technology*, 8(3), 609-612.

Chidankumar, C. S., Chandraju, S., & Nagendraswamy, R. (2009). Impact of distillery spentwash irrigation on the yields of top vegetables (Creepers). *World applied sciences journal*, 6(9), 1270-1273.

Choudhary, A. N., Farooq, M. S., Zeeshan, M., Khan, G., Choudhary, T. K., & Abbas, M. S. (2018). Crop yield and soil characteristics as affected by composts from different organic materials with spent wash. *Adv. Crop Sci. Tech*, 5(2).

Department of Agriculture, Sri Lanka. (2021). Retrieved from <https://doa.gov.lk/hordi-crop-tomato/>

Diangan Jr, M., Perez, T., & Claveria, R. (2008). Analysis of land application as a method of disposal for distillery effluent. *International Journal of Environment and Health*, 2(2), 258-271.

Elevation map for Badulla, Sri Lanka. (2020). Flood Map. Retrieved from <https://www.floodmap.net/Elevation/ElevationMap/?gi=1250615>.

FAO. 2021. World Food and Agriculture – Statistical Yearbook 2021. Rome. Retrieved from <https://doi.org/10.4060/cb4477en>

Fiskesjö, G., 1985. The Allium test as a standard in environmental monitoring. *Hereditas*, 102(1), pp.99-112.

Fiskesjö, G., 1988. Allium test in front of video display units. *Hereditas*, 108(2), pp.239-242.

- Fito, J., Tefera, N., & Van Hulle, S. W. (2019^a). Sugarcane biorefineries wastewater: bioremediation technologies for environmental sustainability. *Chemical and Biological Technologies in Agriculture*, 6(1), 1-13.
- Fito, J., Tefera, N., Kloos, H., & Van Hulle, S. W. (2019^b). Physicochemical properties of the sugar industry and ethanol distillery wastewater and their impact on the environment. *Sugar Tech*, 21(2), 265-277.
- Gana, J.M., Ordóñez, R., Zampini, C., Hidalgo, M., Meoni, S. and Isla, M.I., 2008. Industrial effluents and surface waters genotoxicity and mutagenicity evaluation of a river of Tucuman, Argentina. *Journal of hazardous materials*, 155(3), pp.403-406.
- Garcia, C. F. H., de Souza, R. B., de Souza, C. P., Christofolletti, C. A., & Fontanetti, C. S. (2017). Toxicity of two effluents from agricultural activity: comparing the genotoxicity of sugar cane and orange vinasse. *Ecotoxicology and environmental safety*, 142, 216-221.
- Hand book of organic manure. (2021). Department of Agriculture.
- Hemachandra, C. K., & Pathiratne, A. (2018). Assessing toxicity of two industrial zone effluents reaching Kelani River, Sri Lanka. *Journal of the National Science Foundation of Sri Lanka*, 46(4), 538-546.
- Hemachandra, C.K. and Pathiratne, A., 2015. Assessing toxicity of copper, cadmium and chromium levels relevant to discharge limits of industrial effluents into inland surface waters using common onion, *Allium cepa* bioassay. *Bulletin of Environmental Contamination and Toxicology*, 94(2), pp.199-203.
- Hemavathi, K. L., Kumar, N. H., & Jagannath, S. (2015). Genotoxic effect of distillery effluent on root tip cells of *Allium sativum* L. *Journal of Applied Biology and Biotechnology*, 3(3), 0-4.
- Hussain, I., Raschid, L., Hanjra, M. A., Marikar, F., & Van Der Hoek, W. (2002). Wastewater use in agriculture: Review of impacts and methodological issues in valuing impacts.

Impacts of trade liberalization on the world sugar market. (n.d). Food and Agriculture Organization. <https://www.fao.org/3/x0513e/x0513e06.html>

Jain, R., & Srivastava, S. (2012). Nutrient composition of spent wash and its impact on sugarcane growth and biochemical attributes. *Physiology and Molecular Biology of Plants*, 18(1), 95-99.

Junior, H.M., Da Silva, J., Arenzon, A., Portela, C.S., de Sá Ferreira, I.C.F. and Henriques, J.A.P., 2007. Evaluation of genotoxicity and toxicity of water and sediment samples from a Brazilian stream influenced by tannery industries. *Chemosphere*, 67(6), pp.1211-1217.

Kannangara, D. N. M., & Pathiratne, A. (2015). Toxicity assessment of industrial wastewaters reaching Dandugan Oya, Sri Lanka using a plant based bioassay. *Journal of the National Science Foundation of Sri Lanka*, 43(2).

Karunyal, S., Renuga, G., & Kailash, P. (1994). Effects of tannery effluent on seed germination, leaf area, biomass and mineral content of some plants. *Bioresource Technology*, 47(3), 215-218.

Kaushik, A., Nisha, R., Jagjeeta, K., & Kaushik, C. P. (2005). Impact of long and short term irrigation of a sodic soil with distillery effluent in combination with bioamendments. *Bioresource technology*, 96(17), 1860-1866.

Kharayat, Y. (2012). Distillery wastewater: bioremediation approaches. *Journal of Integrative Environmental Sciences*, 9(2), 69-91.

Koizumi, T. (2003). The Brazilian ethanol programme: impacts on world ethanol and sugar markets and Agriculture, 113.

Kristen, U., 1997. Use of higher plants as screens for toxicity assessment. *Toxicology in vitro*, 11(1), pp.181-191.

Kulasekara, B. R., Weerasinghe, H. A. S., & Ariyawansha, B. D. S. K. Production of Organic Fertilisers by Using Sugarcane Industry By-products of Sri Lanka: A Preliminary Investigation.

- Kumar, V., & Chopra, A. K. (2012). Fertigation effect of distillery effluent on agronomical practices of *Trigonella foenum-graecum* L. (Fenugreek). Environmental Monitoring and Assessment, 184(3), 1207-1219.
- Kumar, V., Shahi, S. K., Ferreira, L. F. R., Bilal, M., Biswas, J. K., & Bulgariu, L. (2021). Detection and characterization of refractory organic and inorganic pollutants discharged in biomethanated distillery effluent and their phytotoxicity, cytotoxicity, and genotoxicity assessment using *Phaseolus aureus* L. and *Allium cepa* L. Environmental Research, 201, 111551.
- Kumari, K., & Phogat, V. K. (2010). Characterization of Spent Wash from Different Distilleries Operating in Haryana and its Utilization as a Source of Liquid Manure in Agriculture. Journal of the Indian Society of Soil Science, 58(3), 347-351.
- Leme, D.M. and Marin-Morales, M.A., 2009. *Allium cepa* test in environmental monitoring: a review on its application. Mutation Research/Reviews in Mutation Research, 682(1), pp.71-81.
- Mahimairaja, S., & Bolan, N. S. (2004). Problems and prospects of agricultural use of distillery spentwash in India. Magnesium, 1715, 2100.
- Mahimairaja, S., & Bolan, N. S. (2004). Problems and prospects of agricultural use of distillery spentwash in India. Magnesium, 1715, 2100.
- Melamane, X. L., Strong, P. J., & Burgess, J. E. (2007). Treatment of wine distillery wastewater: a review with emphasis on anaerobic membrane reactors. South African Journal of Enology and Viticulture, 28(1), 25-36.
- Mikucka, W., & Zielińska, M. (2020). Distillery stillage: characteristics, treatment, and valorization. Applied biochemistry and biotechnology, 192(3), 770-793.
- Mohana, S., Acharya, B. K., & Madamwar, D. (2009). Distillery spent wash: treatment technologies and potential applications. Journal of Hazardous Materials, 163(1), 12-25.

- Oñate, J., Arenas, A., Ruiz, A., Rivera, K., & Peláez, C. (2015). Evaluation of mutagenic and genotoxic activity in vinasses subjected to different treatments. *Water, Air, & Soil Pollution*, 226(5), 1-10.
- Pandey, S. K., Gupta, A. K., & Yunus, M. (2005). Physico-chemical analysis of treated distillery effluent irrigation responses on crop plants pea (*Pisum sativum*) and wheat (*Triticum aestivum*). *Life Science Journal*.
- Pathiratne, A., Hemachandra, C.K. and De Silva, N., 2015. Efficacy of *Allium cepa* test system for screening cytotoxicity and genotoxicity of industrial effluents originated from different industrial activities. *Environmental monitoring and assessment*, 187(12), pp.1-12.
- Pedro-Escher, J., Christofoletti, C. A., Ansoar-Rodríguez, Y., & Fontanetti, C. S. (2016). Sugarcane vinasse, a residue of ethanol industry: toxic, cytotoxic and genotoxic potential using the *Allium cepa* test. *Journal of Environmental Protection*, 7(05), 602.
- Pikaev, A. K., Ponomarev, A. V., Bludenko, A. V., Minin, V. N., & Elizar'eva, L. M. (2001). Combined electron-beam and coagulation purification of molasses distillery slops. Features of the method, technical and economic evaluation of large-scale facility. *Radiation Physics and Chemistry*, 61(1), 81-87.
- Prado, R. D. M., Caione, G., & Campos, C. N. S. (2013). Filter cake and vinasse as fertilizers contributing to conservation agriculture. *Applied and Environmental Soil Science*, 2013.
- Radić, S., Stipaničev, D., Vujičić, V., Rajčić, M. M., Širac, S., & Pevalek-Kozlina, B. (2009). The evaluation of surface and wastewater genotoxicity using the *Allium cepa* test. *Science of the Total Environment*, 408(5), 1228-1233.
- Ramana, S., Biswas, A. K., Kundu, S., Saha, J. K., & Yadava, R. B. R. (2002). Effect of distillery effluent on seed germination in some vegetable crops. *Bioresource technology*, 82(3), 273-275.

- Saliha, B. B., & Manivannan, M. I. (2008). Influence of distillery spentwash on the physico-chemical properties of calcareous sodic soil (Calcic natrustalfs). *Asian Journal of Bio Science*, 3(1), 145-151.
- Satyawali, Y., & Balakrishnan, M. (2008). Wastewater treatment in molasses-based alcohol distilleries for COD and color removal: a review. *Journal of environmental management*, 86(3), 481-497.
- Shinde, P. A., Ukarde, T. M., Pandey, P. H., & Pawar, H. S. (2020). Distillery spent wash: An emerging chemical pool for next generation sustainable distilleries. *Journal of Water Process Engineering*, 36, 101353.
- Siddiqui, A.H., Tabrez, S. and Ahmad, M., 2011. Validation of plant-based bioassays for the toxicity testing of Indian waters. *Environmental monitoring and assessment*, 179(1-4), pp.241-253.
- Srivastava, P. C., Singh, R. K., Srivastava, P., & Shrivastava, M. (2012). Utilization of molasses-based distillery effluent for fertigation of sugarcane. *Biodegradation*, 23(6), 897-905.
- Srivastava, S., & Jain, R. (2010). Effect of distillery spent wash on cytomorphological behaviour of sugarcane settling. *Journal of Environmental Biology*, 31(5), 809.
- Stages of mitosis. n.d. Retrieved from:<https://sciencing.com/5-stages-mitosis-13121.html>
- Sustainable Biofuel for Sri Lanka. 2012. Sustainable Energy Authority. https://rise.esmap.org/data/files/library/srilanka/Renewable%20Energy/RE%206.1%20Sustainable%20Biofuel%20for%20Sri%20Lanka_SLSEA.pdf.
- Tedesco, S., & Laughinghouse IV, H. (2012). Bioindicator of genotoxicity: the Allium cepa test. *Environmental contamination*, 137-156.
- Tewari, P. K., Batra, V. S., & Balakrishnan, M. (2007). Water management initiatives in sugarcane molasses-based distilleries in India. *Resources, Conservation and Recycling*, 52(2), 351-367.

Umair Hassan, M., Aamer, M., Umer Chattha, M., Haiying, T., Khan, I., Seleiman, M. F., & Huang, G. (2021). Sugarcane distillery spent wash (DSW) as a bio-nutrient supplement: a win-win option for sustainable crop production. *Agronomy*, 11(1), 183.

Verma, N., & Dhiman, A. K. A Comparative Study of Effect of Untreated and Anaerobically treated Distillery Effluent on the Seedling Growth of Potato (*Solanum tuberosum* CV G4).

Vujošević, M., Andđelković, S., Savić, G., & Blagojević, J. (2008). Genotoxicity screening of the river Rasina in Serbia using the Allium anaphase–telophase test. *Environmental monitoring and assessment*, 147(1), 75-81.